


Early Detection and Prevention saves lives

- One in every 20 Americans over the Age of 50 has PAD (Peripheral Arterial Disease). PAD Increases the Risk of Heart Attack and Stroke.
- Up to 3% of individuals older than 65 have carotid artery disease. Stroke Causes 1 in every 15 deaths.
- Every year 200,000 people in the US are diagnosed with an abdominal aortic aneurysm(AAA). A ruptured AAA is the 15th leading cause of death in the



Vascular Risk Assessment and Screening



- ✉ contact@jrkmicals.com
- 📍 2633 Superior Drive
NW Suite 100
Rochester MN 55901
- ☎ 507-322-6900

We offer Screening for

1. Peripheral Arterial Disease
2. Carotid Artery Disease
3. Abdominal Aortic Aneurysm

All Three scans for \$ 149 Only





Come Prepared for Easy Screening

- Wear two piece loose fitting clothing, allows for easy scanning.
- Avoid tights or pantyhose.
- Aneurysm screening need Fast for minimum 4 hours.
- Do not eat for 8-12 hours if checking for fasting glucose or lipid, can drink water or take medications.
- Do not use oils, creams or lotion on the skin, can interfere with scan.
- If you are a diabetic please consult your doctor for instructions on medications.
- Take medications as prescribed.

Vascular Screening and Vascular Risk Assessment can facilitate

- **Early detection of peripheral arterial disease.**
- **Early detection of stroke risk** due to Blockage in the Carotid Artery.
- **Early detection of an aneurysm**, particularly if you are a current or past smoker and have family history of an aneurysm.

Benefits of Vascular Screening and Vascular Risk Assessment

- Early detection allows early intervention to prevent disease progression
- Most Screenings are affordable, **does not need insurance and takes less than an hour to complete.**
- Results are immediately available and also at your request can be mailed to your primary care provider.